

Guoyong Fu

*Princeton Plasma Physics Laboratory
Princeton, NJ*

Nonlinear Hybrid Simulations of Multiple Energetic Particle driven Alfvén Modes in Toroidal Plasmas

Understanding of nonlinear behavior of energetic particle-driven instabilities in tokamaks is of fundamental importance for burning plasmas. Here we report recent advances in self-consistent nonlinear simulations of fast beam ion-driven Alfvén modes in NSTX. In the hybrid model, the thermal electrons and ions are treated as an ideal fluid while the energetic species is described by either drift-kinetic equation or gyrokinetic equation. The effects of energetic particles are coupled to the MHD equations via the stress tensor term in the momentum equation. The hybrid code has been recently applied to study nonlinear dynamics of fishbone instability [2]. The code was also used to simulate nonlinear evolution of a single beam-driven TAE mode in NSTX. The result showed a weak frequency chirping about 20% consistent with experimental measurement [3]. In this work, we use the M3D code to simulate beam ion driven Alfvén modes in NSTX plasmas with multiple unstable Alfvén modes. It is found that mode saturation level of each mode can be enhanced significantly by presence of other unstable modes indicating strong nonlinear interaction between different modes. It is also found that a linearly stable $n=2$ mode can be nonlinearly driven by an $n=1$ mode at significant mode amplitude.

[1] W. Park, E.V. Belova, G.Y. Fu et al., *Phy. Plasmas* 6, 1796 (1999)

[2] Fu, GY, Park, W, Strauss HR et al. *PHYSICS OF PLASMAS* 13, 052517 (2006)

[3] Fu GY, Breslau J, Fredrickson E et al. ,*Proceedings of 2004 IAEA Fusion Energy Conference, Vilamoura, Portugal, Paper TH/P4-38.*
